

## What can I do to reduce LPR?

There are things you can do to reduce the likelihood of LPR and acid indigestion:

- **LOSE WEIGHT** and eat a healthy diet
- **LIMIT** your intake of fried foods, spicy foods, chocolate and citrus juices
- **GIVE UP** smoking
- **RAISE THE HEAD OF YOUR BED** 4-6 inches (place blocks under the legs/base of the bed at the head end)
- **LEAVE 3 HOURS** between eating and lying down
- **BEND AT THE KNEES** when you pick things up
- Wear **LOOSE CLOTHING** around your waist
- **CHEW GUM** after meals
- **EAT SMALLER MEALS** more regularly
- **EAT SLOWLY**, chewing each mouthful well
- **AVOID FIZZY DRINKS and CAFFEINE** – drink water or herbal teas instead
- **DRINK LESS ALCOHOL**, especially before bed

## Will I always need treatment for LPR?

Usually LPR improves with the appropriate treatment but you need to:

- Follow the recommendations you are given on lifestyle changes
- Make the dietary changes that are recommended
- Take your medication regularly as prescribed by your doctor
- Remember, any decision to stop treatment should be made with your doctor's knowledge and consent.

For more information about laryngo-pharyngeal reflux see the BVA website:

[www.britishvoiceassociation.org.uk](http://www.britishvoiceassociation.org.uk)

**Acknowledgements:** The BVA are grateful to the Speech and Language Therapy and Ear, Nose and Throat teams at Lewisham Healthcare NHS Trust, London UK for allowing us to adapt their material in the production of this leaflet and to Tomm Coles, Nutritionist, John Rubin, ENT Surgeon/Laryngologist and Tom Harris, ENT Surgeon/Laryngologist for advice.

330 Gray's Inn Road, London WC1X 8EE

Tel: +44 (0)300 123 2773

email: [adminstrator@britishvoiceassociation.org.uk](mailto:adminstrator@britishvoiceassociation.org.uk)



# REFLUX AND YOUR VOICE

## WHAT IS ACID REFLUX?

**Stomach juices are made up of strong digestive acids and enzymes. It is not uncommon for these juices to travel from the stomach upwards into the gullet (oesophagus) causing the symptoms of indigestion (heartburn). This is usually referred to as 'gastro-oesophageal' reflux (GOR).**

Sometimes, small amounts of these stomach juices can reach as far up as the throat (pharynx) and voice box (larynx). This is known as laryngo-pharyngeal reflux (LPR) and is sometimes called 'silent reflux' as many people do not experience symptoms of heartburn. Acid reflux can occur during the day or night, even if a person hasn't eaten anything.



A healthy larynx

*Photo: Tom Harris*

This leaflet is intended for guidance purposes only and is in no way intended to replace professional clinical advice by a qualified practitioner.

## What are the symptoms?

These may vary but some of the most common symptoms of LPR are:

- A sensation of food sticking or a feeling of a lump in the throat
- A hoarse, tight or 'croaky' voice
- Frequent throat clearing
- Difficulty swallowing (especially tablets or solid foods)
- A sore, dry and sensitive throat
- Occasional unpleasant "acid" or "bilious" taste at the back of the mouth
- A feeling that too much mucus/phlegm is collecting in the throat
- Sudden coughing or choking spasms at night
- Chronic cough
- Excessive burping, particularly during the day

Boggy swelling (oedema) at the back of the larynx

Reddening (Erythema) around structures at back of the larynx




Photo: Tony Aymat

## How is LPR diagnosed?

LPR is usually diagnosed by examination of your throat and voice box by an Ear, Nose and Throat (ENT) doctor. The ENT doctor or the Speech and Language Therapist (SLT) specialising in voice disorders will ask you to describe your symptoms. They may also ask you to fill in a questionnaire to rate how your symptoms affect you.

If your symptoms are severe your doctor may send you for further tests to measure the level of acid in your throat and gullet and, if necessary, refer you on to a gastroenterologist (a doctor specialising in stomach and digestive problems).

## What causes LPR?

Often it is not clear what causes LPR. There may be a number of underlying causes, such as problems with the stomach, the gullet or the sphincter muscle that separates the two. But, in many cases, lifestyle factors such as being overweight, smoking and consuming too much alcohol and caffeine are probable causes. Dietary factors may also play an important role with spicy foods, fatty foods, chocolate and citrus juice contributing to symptoms of LPR in some individuals.

## How is LPR treated?

Treatment is usually planned on an individual basis and your doctor or SLT will suggest the appropriate treatment for you. There are several treatments that are effective:

- Making lifestyle and dietary changes
- Taking medications to reduce stomach acid
  - Non-prescription antacids/alginate preparations
  - Acid-blocking tablets called proton pump inhibitors. (These must be prescribed by your doctor)
- Various surgical options might be recommended to tighten the valve between the stomach and the gullet when medication has not been effective.

Most people with LPR report improvement in their symptoms within 2-3 months of treatment but it may take 6 months or more for the throat and voice symptoms to resolve. Stopping reflux medications suddenly can result in increased LPR, so most doctors recommend a 'step down' plan. This will be explained by your doctor and it is important that you follow your doctor's advice.